

## TAXONOMIC ACCOUNT OF EUGLENOIDS IN SELECTED SITES OF KARUVANNUR RIVER, KERALA, INDIA

Devikrishna C.S.<sup>1\*</sup>, Tessy Paul P.<sup>2</sup> and Mohamed Nasser K.M.<sup>1</sup>

<sup>1</sup>. Research & PG Department of Botany, M.E.S. Asmabi College, P.Vemballur, Kerala – 680671

<sup>2</sup>. Department of Botany, Christ College (Autonomous), Irinjalakuda, Kerala- 680 125

<sup>1 & 2</sup> Affiliated to University of Calicut

\*Corresponding author: [csdevi777@gmail.com](mailto:csdevi777@gmail.com)

### ABSTRACT

The Euglenophytes are one of the important groups of phytoplankton which is primarily freshwater inhabitants and are unicellular flagellates. The present paper is an attempt to explore the euglenoid diversity and to make a taxonomic account on it, from the selected sites of Karuvannur River, Thrissur District, Kerala. Karuvannur River is one of the major freshwater sources of Thrissur district, which is flowing through the famous Kole lands of Thrissur. The study was carried out from June 2017 to May 2018. During the period of study, 25 species of euglenophyceae members were identified belonging to five genera, namely, *Euglena*, *Lepocinclis*, *Phacus*, *Trachelomonas* and *Strombomonas*.

**Keywords:** Euglenoid diversity, Karuvannur River, *Euglena*, *Lepocinclis*, *Phacus*, *Trachelomonas*, *Strombomonas*.

### INTRODUCTION

Algae play a vital role in maintaining the food chain since they are the primary producers. These lower plants are used as feed, fodder, fertilizer, medicines and even as a source of restriction endonucleases (John and Francis, 2012). Also they have a noticeable role in the ecological aspects, markedly as pollution indicators (Palmer, 1969). Euglenoids are unicellular flagellates primarily occur in freshwater habitat. Although a few are sedentary, most forms are motile with one or two flagella and many of them are highly metabolic, even with an amoeboid type of motility (Prescott, 1982). They are often predominant in eutrophic waters including high organic and inorganic contents (Rahman *et al*, 2014).

Considerable work has been carried out in India about systematic survey, distribution and seasonal occurrence of euglenoids (Suxena, 1955; Munavar, 1972; Pandey and Pandey, 1980; Ratha *et al*, 2006). Philipose (1982, 1984, 1988) has done a very extensive work on euglenophytes throughout India.

Very few reports are available on the euglenoid diversity, from the freshwater habitats of Kerala. Shaji and Patel (1991), Shajiet *al* (1995), John and Francis (2012) and Arulmurugan *et al* (2010) have described euglenoids from freshwaters of Kerala. This is an attempt to explore the euglenoid diversity and to make a taxonomic account on it,

from the selected sites of Karuvannur River, Thrissur District, Kerala.

### MATERIALS AND METHODS

Karuvannur River is one of the major freshwater sources of Thrissur district, which is flowing through the famous Kole lands of Thrissur, with its origin at Pumalai hills in Chimmony Wildlife sanctuary. It has two main tributaries, Kurumali River and Manali River, which confluence at Palakkadavu, before Arattupuzha to form the Karuvannur River. For this study ten sampling sites were selected, which include the beginning and end region of the river, viz, Chimmony (S1), Peechi (S2), Kurumali (S3), Puthur (S4), Manali (S5), Palakkadavu (S6), Arattupuzha (S7), Karuvannur (S8), Thriprayar (S9) and Enammavu (S10). The surface water samples were collected from the sites at monthly intervals from June 2017 to May 2018. Collected samples were preserved by adding 4% formalin (APHA, 1998). Each sample was mounted on the glass slide using glycerine and observed thoroughly under research microscope for the taxonomic analysis. Identification of algal forms was made with the help of relevant and available floras and literatures (Prescott, 1982; Philipose, 1982, 1984, 1988; Wolowski, 1998). Photomicrographs of algal taxa taken with digital camera attached to the research microscope.

### RESULTS AND DISCUSSION

During the period of study, 25 species of euglenoids were identified which belongs to five genera, namely, *Euglena*, *Lepocinclis*, *Phacus*, *Trachelomonas* and *Strombomonas* from the selected sites of Karuvannur River.

#### Taxonomic description:

Genus: *Euglena* Ehrenberg

#### 1. *Euglena acus* Ehr.

Prescott, 1982. p. 390, pl. 85, fig. 28.

Cells elongate, spindle shaped, 140-150 µm long, 10-14.6 µm broad, produced posteriorly into a long, fine tapering point, narrowed and truncate at the anterior end. [Col. Site: S5, S10]

#### 2. *Euglena charkowiensis* Swirensko

Islam and Irfanullah, 2005. pl. 4, fig. 47.

Cells 140–153.0 µm long, 14.6–20.9 µm broad, posterior end with a straight tail-piece. [Col. Site: S3, S5]

**3. *Euglena oblonga* Schmitz.**

Khondkeret *et al.*, 2008. Figs. 15a–c.

Cells elongated, ovoid-spindle, 22–25.5  $\mu\text{m}$  broad, 72.2–82  $\mu\text{m}$  long, anterior narrowed to a rounded end, posterior suddenly narrowed to a blunt end. [Col. Site: S2, S8, S10]

**4. *Euglena proxima* Dangeard**

Prescott, 1982. p. 394, pl. 85, fig. 25.

Cells fusiform, 48.0–57.3  $\mu\text{m}$  long, 14.5–23.3  $\mu\text{m}$  broad, cells, narrowed posteriorly to blunt hyaline tip. [Col. Site: S7, S8, S10]

**5. *Euglena spathiryncha* Skuja**

Khondkeret *et al.*, 2008. Figs. 19a–c.

Cells 109–124.5  $\mu\text{m}$  long, 21.4–35  $\mu\text{m}$  broad, cells elongated, spindle, mid-region bulged out, posterior gradually narrowed to a sharp pointed end [Col. Site: S10]

**6. *Euglena sp.1***

Wolowski, 1998.

Cells 61.2–67.0  $\mu\text{m}$  long, 20.7–23.5  $\mu\text{m}$  broad, cell slightly extending and rounded at the anterior end, narrowing to short tail-piece at the posterior end. [Col. Site: S7]

Genus: *Phacus* Dujardin

**7. *Phacusanacoelus* Stokes**

Prescott, 1982. p. 397, pl. 87, figs. 7, 8 and pl. 88, fig. 11.

Cells broadly ovoid, posterior end abruptly narrowed to a short caudus, cells 41.2–56.6  $\mu\text{m}$  long, 33.2–41.2  $\mu\text{m}$  broad. [Col. Site: S7]

**8. *Phacushelikoides* Pochmann**

Prescott, 1982. p. 400, pl. 87, fig. 9.

Cell spirally twisted, elongate pyriform, markedly broad in the anterior third, and with a straight hyaline tapering tail at the posterior end, 49.2–52.5  $\mu\text{m}$  broad, 112–128.3  $\mu\text{m}$  long. [Col. Site: S5]

**9. *Phacus sp. 1***

Philipose, 1984.

Cells elongate-ellipsoid, posterior end with a short straight tail, cells 18.3–20.5  $\mu\text{m}$  broad, 28.8–31.0  $\mu\text{m}$  long, periplast with longitudinal rows of small forwardly pointed spines. [Col. Site: S10]

**10. *Phacus sp.2***

Wolowski, 1998.

Cells 50.2–55.7  $\mu\text{m}$  long, 36.05–38.0  $\mu\text{m}$  wide, oval, corrugated at the rim, with a long curved cauda at the posterior end. [Col. Site: S5]

Genus: *Lepocinclis* Perty

**11. *Lepocinclis acuta* Prescott**

Prescott, 1982. p. 405, pl. 89, figs. 8, 9.

Cells ovoid-pyriform, tapering posteriorly to a long, sharply pointed caudus, slightly narrowed anteriorly, 15–18.1  $\mu\text{m}$  in diameter, 34–38.8  $\mu\text{m}$  long. [Col. Site: S1]

**12. *Lepocinclis acicularis* Francé**

Wolowski, 1998, p.68, figs. 218, 219

Cell 18.0–25.0  $\mu\text{m}$  long, 8.0–8.5  $\mu\text{m}$  wide, fusiform, pellicle with a few spiral striae. [Col. Site: S5, S7]

**13. *Lepocinclis fusiformis* (Carter) Lemm.**

Prescott, 1982. p. 406, pl. 89, fig. 1-4.

Cells broadly pyriform, slightly produced posteriorly to form a blunt basal point, 17–25  $\mu\text{m}$  in diameter, 33–36  $\mu\text{m}$  long. [Col. Site: S7, S10]

**14. *Lepocinclis ovum* (Ehrenb.) Lemm.**

Prescott, 1982. p. 407, pl. 89, figs. 5, 6.

Cells broadly ovate with a short blunt caudus, 15.0–22.0  $\mu\text{m}$  wide, 26.0–30.5  $\mu\text{m}$  long, rounded both anteriorly and posteriorly. [Col. Site: S5, S7, S10]

Genus: *Trachelomonas* Ehrenberg

**15. *Trachelomonas abrupta* (Swir.) Defl.**

Prescott, 1982. p. 410, pl. 83, figs. 18, 19.

Lorica cylindrical, 16.0–17.5  $\mu\text{m}$  broad, 25.5–28.0  $\mu\text{m}$  long, wall coarsely punctate, covered by small blunt spines. [Col. Site: S3, S7, S10]

**16. *Trachelomonas armata* (Ehr.) Stein. var. *longispina* (Playf.) Defl.**

Prescott, 1982. p. 411, pl. 83, fig. 27.

Test 28.2–30.6  $\mu\text{m}$  broad, 32.0–41.5  $\mu\text{m}$  long, broadly obovate, flagellum aperture with a circle of erect spines at the margin, anterior margin with short and posterior with both short and long stout spines. [Col. Site: S5, S7, S10]

**17. *Trachelomonas dubia* (Swiremend) Defl.**

Prescott, 1982. p. 412, pl. 85, figs. 1, 2

Lorica cylindrical, smooth, anterior end abruptly narrowed to form a short cylindrical neck, 10.5–12.6  $\mu\text{m}$  broad, 23.5–28.0  $\mu\text{m}$  long. [Col. Site: S3]

**18. *Trachelomonas hispidia* (Perty) Stein. var. *duplex* Playf.**

Prescott, 1955. pl. 2, fig. 8.

Test ovate, cells 24.5–27.0  $\mu\text{m}$  long, 17.5–20.4  $\mu\text{m}$  in diameter, narrowed anteriorly, flagellum aperture slightly raised, wall uniformly covered with minute sharp spines. [Col. Site: S3]

**19. *Trachelomonas lacustris* Drez.**

Prescott, 1982. p. 415, pl. 83, figs. 14, 15.

Lorica 13.0–15.0  $\mu\text{m}$  broad, 22.0–24.0  $\mu\text{m}$  long, cylindrical, flagellum aperture with slightly raised rim, wall densely punctate. [Col. Site: S9, S10]

**20. *Trachelomonas planctonica* fo. *ornata* Skvortzov**

Wolowski, 1998. p.62, Fig. 199.

Test 17.0–21.5  $\mu\text{m}$  wide, 20.0–32.0  $\mu\text{m}$  long, elliptical, verrucose and covered with pores, collar with irregular rim. [Col. Site: S6, S10]

**21. *Trachelomonas robusta* Swirenko**

Prescott, 1982. p. 416, pl. 83, fig. 29.

Test subglobose, 15.5–17.2  $\mu\text{m}$  broad, 20.3–25.2  $\mu\text{m}$  long, flagellum aperture with a thickened rim, wall dark brown, evenly beset with short sharp spines. [Col. Site: S3, S4, S5]

**22. *Trachelomonas superba* (Swir.) Defl. var. *swirenkiana* Defl.**

Prescott, 1982. p. 418, pl. 84, figs. 8, 9 and pl. 83, fig. 34.

Test subglobose, 29.5–32  $\mu\text{m}$  broad, 39.3–41.5  $\mu\text{m}$  long, flagellum aperture low ring like collar, anterior and posterior wall spiny, posterior spines longer and stouter than anterior region. [Col. Site: S5]

**23. *Trachelomonasvolvocina* Ehr.**

Prescott, 1982. p. 419, pl. 83, figs. 1, 7, 8.

Lorica globose, smooth, 14.5–20.4 µm in diameter, collar usually absent. [Col. Site: S1, S2, S3, S5, S7, S8, S9, S10]

**24. *Trachelomonas* sp.1**

Philipose, 1988.

Lorica spindle-shaped, cells 17.0–20.0 µm broad, 50.0–58.8 µm long. [Col. Site: S5, S10].

Genus: *Strombomonas* Deflandre

**25. *Strombomonasfluviatilis* (Lemm.) Defl.**

Islam and Irfanullah, 2005. pl. 3, fig. 44

Lorica fusiform, cells 21.5–26.0 µm broad, 46.0–50.6 µm long, anterior end narrowed, with slightly widened cylindrical neck; narrowed at the posterior end into conical appendix. [Col. Site: S9].

Out of the 25 species, *Trachelomonas* represents more number with 10 species, followed by *Euglena* with 6 species in it. *Lepocinclis* and *Phacus* have 4 species each. *Strombomonas* is represented by single species *S. fluviatilis*. It is *Trachelomonasvolvocina* seen to be present in most of the selected sites.

**CONCLUSION**

This study was to explore the euglenoid diversity of some selected sites from Karuvannur River, Thrissur District. 25 species of euglenoids were observed in the study period which belongs to 5 genera, *Euglena* (6 spp.), *Phacus* (4 spp.), *Lepocinclis* (4 spp.), *Trachelomonas* (10 spp.) and *Strombomonas* (1 spp.).

**ACKNOWLEDGEMENTS**

We wish to thank Principal, MES Asmabi College and Head, Dept. of Botany for the permission to use laboratory and various helps during the study.

**REFERENCES**

1. APHA. (1998). *Standard methods for the examination of water and waste water*. 20<sup>th</sup> edn. American Public Health Association, Washington DC.
2. Arulmurugan, P., Nagaraj, S. & Anand, N. (2010). Biodiversity of freshwater algae from temple tanks of Kerala. *Rec. Res. Sci. Tech.* 2: 58-72.
3. Islam, A.K.M., Nurul & Irfanullah, H.M. (2005). Hydrobiological studies within the tea gardens at Srimangal, Bangladesh. II. Algal flora (excluding Chlorophyceae). *Bangladesh J. Plant Taxon.*, 12(1): 33–52.
4. John, J. and Francis M.S. (2012). *An illustrated algal flora of Kerala*. Vol. I. Idukki district. Pranatha books, Kochi. 281 pp.
5. Khondker, M., Bhuiyan, R.A., Yeasmin, J., Alam, M., Sack, R.B., Huq, A. and Colwell, R.R. (2008). New records of phytoplankton for Bangladesh. 5. *Euglena*, *Euglenocapsa*. *Bangladesh J. Plant Taxon.*, 15(1): 39–46.
6. Munawar, M. (1972). Ecological studies of Euglenineae in certain polluted and unpolluted environments. *Hydrobiologia*, 39: 207-320.
7. Palmer, C.M. (1969). A composite rating of algae tolerating organic pollution. *J. Phycol.*, 5: 78–82.
8. Pandey, U. C. and Pandey, D. C. (1980). Freshwater Euglenineae from Allahabad. *Kanpur Univ. Res. J.* 1: 247-251.
9. Philipose, M.T. (1982). Contributions to our knowledge of Indian algae-III- Euglenineae-Part-I. The genus *Euglena* Ehrenberg. *Proc. Indian Acad. Sci. (Plant Sci.)*, 91(6): 551-599.
10. Philipose, M.T. (1984). Contributions to our knowledge of Indian algae-III- Euglenineae-Part 2. *Proc. Indian Acad. Sci. (Plant Sci.)*, 93(5): 503-552.
11. Philipose, M.T. (1988). Contributions to our knowledge of Indian algae-III- Euglenineae-Part 3. The genera *Trachelomonas* Ehrenberg and *Strombomonas* Deflandre. *Proc. Indian Acad. Sci. (Plant Sci.)*, 98(5): 317-394.
12. Prescott, G.W. (1955). Algae of the Panama canal and its tributaries I. Flagellated organisms. *The Ohio Journal of Science*, 55(2): 99–121.
13. Prescott, G.W. (1982). *Algae of the Western Great Lakes Area, With an illustrated key to the Genera of Desmids and Freshwater Diatoms*. Koenigstein Otto Koeltz. 977 pp.
14. Rahman, M. M., Ghosh, J. K. and Islam, M. S. (2014). Relationships of euglenophytes bloom to environmental factors in the fish ponds at Rajshahi, Bangladesh. *Journal of Agriculture and Veterinary Science*, 7(1): 45-52.
15. Ratha, S. K., Jena, M. and Adhikary, S. P. (2006). Euglenophytes from Orissa State, East Coast of India. *Algae*, 21(1): 61-73.
16. Shaji, C. and Patel, R. J. (1991). Contributions to Euglenoids of Kerala, India. *Phykos*, 30(1&2): 109-114.
17. Shaji, C., Sindhu, P. and Panikkar, M. V. N. (1995). Contributions to Euglenoids of Kerala, India - II. *J. Econ. Tax. Bot.*, 19 (2): 269-272.
18. Suxena, M. R. 1955. Freshwater Euglenineae from Hyderabad, India. *J. Indian bot. Soc.* 34(4): 429-450.
19. Wołowski, K. (1998). Taxonomic and environmental study on euglenophytes of the Kraków Częstochowa upland (southern Poland). *Fragm. Florist. Geobot. Suppl.* 6: 1–192.